

A complete asymptotic expansion of power means

Ulrich Abel^{a,*}, Mircea Ivan^b

^a *Fachbereich MND, Fachhochschule Gießen-Friedberg, University of Applied Sciences, Wilhelm-Leuschner-Straße 13, 61169 Friedberg, Germany*

^b *Department of Mathematics, Technical University Cluj-Napoca, Str. C. Daicoviciu 15, 400020 Cluj-Napoca, Romania*

Received 31 December 2005

Available online 28 February 2006

Submitted by William F. Ames

Abstract

The complete asymptotic expansion of power means in terms of Bell polynomials is obtained. Some results recently obtained by M. Bjelica are generalized.

© 2006 Elsevier Inc. All rights reserved.

Keywords: Asymptotic expansions; Means

1. Introduction and results

Let $x_i > 0$, $w_i \geq 0$, $1 \leq i \leq n$, $\sum_{i=1}^n w_i = 1$, $\mathbf{x} = (x_1, \dots, x_n)$.

The power mean of order p is defined as

$$M_p(\mathbf{x}) := \left(\sum_{i=1}^n w_i x_i^p \right)^{1/p} \quad (p \in \mathbb{R} \setminus \{0\}) \quad (1)$$

and in the case $p = 0$ as

$$M_0(\mathbf{x}) := \prod_{i=1}^n x_i^{w_i}.$$

* Corresponding author.

E-mail addresses: ulrich.abel@mnd.fh-friedberg.de (U. Abel), mircea.ivan@math.utcluj.ro (M. Ivan).

It is easy to show that

$$M_0(\mathbf{x}) = \lim_{p \rightarrow 0} M_p(\mathbf{x}).$$

Throughout the paper let $\mathbf{a} = (a, \dots, a)$ be a point with equal coordinates.

Recently M. Bjelica has studied the asymptotic behaviour of $M_p(\mathbf{x} + \mathbf{a})$ as a tends to infinity. He obtained the following asymptotic relation.

Theorem A. (See [1].) *For each $p \in \mathbb{R} \setminus \{0\}$, the power mean of order p satisfies the asymptotic formula*

$$M_p(\mathbf{x} + \mathbf{a}) = M_1(\mathbf{x} + \mathbf{a}) + (p - 1)(M_2(\mathbf{x} + \mathbf{a}) - M_1(\mathbf{x} + \mathbf{a})) \\ + o(M_2(\mathbf{x} + \mathbf{a}) - M_1(\mathbf{x} + \mathbf{a})) \quad (a \rightarrow +\infty).$$

The purpose of this note is to give a complete asymptotic expansion of $M_p(\mathbf{x} + \mathbf{a})$ in the form

$$M_p(\mathbf{x} + \mathbf{a}) = a + \sum_{k=0}^{\infty} \frac{c_k(p)}{a^k} \quad (a \rightarrow \infty), \quad (2)$$

where the coefficients $c_k(p)$ are independent of a . We express the coefficients $c_k(p)$ in terms of partial and complete Bell polynomials.

Recall that the (exponential) *partial Bell polynomials* are the polynomials $\mathbf{B}_{m,j}[y_i] := \mathbf{B}_{m,j}(y_1, \dots, y_{m-j+1})$ in an infinite number of variables y_1, y_2, \dots , defined by the formal series expansion [2, p. 133, Eq. [3a']]:

$$\frac{1}{j!} \left(\sum_{k=1}^{\infty} y_k \frac{t^k}{k!} \right)^j = \sum_{m=j}^{\infty} \mathbf{B}_{m,j}[y_i] \frac{t^m}{m!}, \quad j = 1, 2, \dots \quad (3)$$

The (exponential) *complete Bell polynomials* $Y_m[y_i] = Y_m(y_1, y_2, \dots, y_n)$ (see, e.g., [2, p. 134, Eq. [3b]]) are defined by

$$\exp \left(\sum_{j=1}^{\infty} y_j \frac{t^j}{j!} \right) = 1 + \sum_{m=1}^{\infty} Y_m[y_i] \frac{t^m}{m!}, \quad (4)$$

in other words,

$$Y_m[y_i] = \sum_{j=1}^m \mathbf{B}_{m,j}[y_i]. \quad (5)$$

Theorem 1. *For each $p \in \mathbb{R}$, the power means $M_p(\mathbf{x} + \mathbf{a})$ possess the complete asymptotic expansion (2). In the case $p \neq 0$ the coefficients are given by*

$$c_k(p) = \frac{1}{(k+1)!} \sum_{j=1}^{k+1} j! \binom{1/p}{j} \mathbf{B}_{k+1,j} \left[i! \binom{p}{i} M_i^i(\mathbf{x}) \right].$$

In the case $p = 0$ the coefficients are given by

$$c_k(0) = \frac{(-1)^{k+1}}{(k+1)!} \mathbf{Y}_{k+1} [-(i-1)! M_i^i(\mathbf{x})].$$

Remark 2. Note that the series (2) is not only a complete asymptotic expansion but also a convergent power-series expansion in a^{-1} converging for all $a > \max_{1 \leq i \leq n} |x_i|$.

The partial Bell polynomials possess the exact expression (see, e.g., [2, p. 134, Theorem A, Eq. [3d]])

$$B_{m,j}[y_i] = \sum m! \prod_{i=1}^{m-j+1} \left(\frac{1}{k_i!} \left(\frac{y_i}{i!} \right)^{k_i} \right), \quad (6)$$

where the summation takes place over all integers $k_1, k_2, \dots \geq 0$, such that

$$k_1 + 2k_2 + 3k_3 + \dots = m \quad \text{and} \quad k_1 + k_2 + k_3 + \dots = j.$$

Taking advantage of this fact we immediately obtain an explicit form of the coefficients $c_k(p)$.

Corollary 3. For each $p \in \mathbb{R}$, the power means $M_p(\mathbf{x} + \mathbf{a})$ possess the complete asymptotic expansion (2). In the case $p \neq 0$ the coefficients are given by

$$c_k(p) = \sum_{j=1}^{k+1} j! \binom{1/p}{j} \sum \prod_{i=1}^{k-j+2} \left(\frac{1}{k_i!} \left[\binom{p}{i} M_i^i(\mathbf{x}) \right]^{k_i} \right)$$

and the summation in the latter sum runs over all integers $k_1, k_2, \dots \geq 0$, such that

$$k_1 + 2k_2 + 3k_3 + \dots = k + 1 \quad \text{and} \quad k_1 + k_2 + k_3 + \dots = j.$$

In the case $p = 0$ the coefficients are given by

$$c_k(0) = (-1)^{k+1} \sum \prod_{i=1}^{k+1} \left(\frac{1}{k_i!} \left[\frac{-1}{i} M_i^i(\mathbf{x}) \right]^{k_i} \right),$$

where the summation runs over all integers $k_1, k_2, \dots \geq 0$ such that

$$k_1 + 2k_2 + 3k_3 + \dots = k + 1.$$

Remark 4. An easy calculation shows that

$$\lim_{p \rightarrow 0} c_k(p) = c_k(0) \quad (k = 0, 1, \dots).$$

For the convenience of the reader we give some explicit expressions of the coefficients $c_k(p)$ which are valid for all $p \in \mathbb{R}$:

$$\begin{aligned} c_0(p) &= M_1(\mathbf{x}), \\ c_1(p) &= \frac{p-1}{2} [M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})], \\ c_2(p) &= \frac{p-1}{3!} [(p-2)M_3^3(\mathbf{x}) - 3(p-1)M_1(\mathbf{x})M_2^2(\mathbf{x}) + (2p-1)M_1^3(\mathbf{x})], \\ c_3(p) &= \frac{p-1}{4!} [(p-2)(p-3)M_4^4(\mathbf{x}) - 4(p-1)(p-2)M_1(\mathbf{x})M_3^3(\mathbf{x}) \\ &\quad - 3(p-1)^2M_2^4(\mathbf{x}) + 6(2p-1)(p-1)M_1^2(\mathbf{x})M_2^2(\mathbf{x}) \\ &\quad - (2p-1)(3p-1)M_1^4(\mathbf{x})]. \end{aligned}$$

We remark that the table of partial Bell polynomials in [2, p. 307] may be helpful for the calculation of further coefficients.

As an immediate corollary of Theorem 1 we obtain an extension of the asymptotic linearity (cf. [1, Eq. (1)]), i.e., of the relation

$$\lim_{\mathbf{x} \rightarrow \mathbf{a}} \frac{M_p(\mathbf{x}) - M_q(\mathbf{x})}{M_r(\mathbf{x}) - M_s(\mathbf{x})} = \frac{p - q}{r - s} \quad (r \neq s).$$

Theorem 5. Let \mathbf{a} be a point with equal coordinates. Furthermore, let \mathbf{x} be a point having not all coordinates equal. For all p, q, r, s with $r \neq s$, we have a complete asymptotic expansion

$$\frac{M_p(\mathbf{a} + t\mathbf{x}) - M_q(\mathbf{a} + t\mathbf{x})}{M_r(\mathbf{a} + t\mathbf{x}) - M_s(\mathbf{a} + t\mathbf{x})} = \sum_{k=0}^{\infty} q_k(p, q, r, s) \left(\frac{t}{a}\right)^k \quad (t \rightarrow 0),$$

where the coefficients $q_k(p, q, r, s)$ are independent of t and a . The first two coefficients are given by

$$q_0(p, q, r, s) = \frac{p - q}{r - s},$$

$$q_1(p, q, r, s) = \frac{1}{3} \frac{p - q}{r - s} (p + q - r - s) \frac{M_3^3(\mathbf{x}) - 3M_1(\mathbf{x})M_2^2(\mathbf{x}) + 2M_1^3(\mathbf{x})}{M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})}.$$

Remark 6. Note that the denominator $M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})$ is different from zero since the power mean $M_p(\mathbf{x})$ is a strictly increasing function of the real variable p unless \mathbf{x} has equal coordinates.

Remark 7. As a special case we obtain an asymptotic relation which can be written in the form

$$\lim_{t \rightarrow 0} \frac{1}{t} \left(\frac{M_p(\mathbf{a} + t\mathbf{x}) - M_q(\mathbf{a} + t\mathbf{x})}{M_r(\mathbf{a} + t\mathbf{x}) - M_s(\mathbf{a} + t\mathbf{x})} - \frac{p - q}{r - s} \right) = \frac{p + q - r - s}{3a} \frac{p - q}{r - s} \frac{M_3^3(\mathbf{x}) - 3M_1(\mathbf{x})M_2^2(\mathbf{x}) + 2M_1^3(\mathbf{x})}{M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})}.$$

Finally, we note that our results are valid for a more general definition of the power mean. Let μ be a positive Borel measure satisfying $\int_{\mathbb{R}} d\mu(x) = 1$. For each positive and μ -integrable function f we put

$$M_p(f) := \begin{cases} \left(\int_{\mathbb{R}} f^p(x) d\mu(x) \right)^{1/p} & (p \in \mathbb{R} \setminus \{0\}), \\ \exp\left(\int_{\mathbb{R}} \log f(x) dx\right) & (p = 0), \end{cases} \quad (7)$$

provided that the integrals exist (cf. [4, p. 64, Eq. (7)], [3, Eq. (1.2)]). It is obvious that definition (1) is a special case of (7).

The complete asymptotic expansion (2) then reads

$$M_p(f + ae_0) = a + \sum_{k=0}^{\infty} \frac{c_k(p)}{a^k} \quad (a \rightarrow \infty),$$

where e_0 denotes the function given by $e_0(x) = 1$ on \mathbb{R} .

The proof of the analogue of Theorem 1 is completely similar. Assume that, in addition, f is bounded. Then in the case $p \neq 0$, for $a > 0$, there holds

$$\begin{aligned}
M_p(f + ae_0) &= aM_p(a^{-1}f + e_0) = a \left(\int_{\mathbb{R}} (1 + a^{-1}f(x))^p d\mu(x) \right)^{1/p} \\
&= a \left(1 + \sum_{k=1}^{\infty} \binom{p}{k} \int_{\mathbb{R}} (a^{-1}f(x))^k d\mu(x) \right)^{1/p} \\
&= a \left(1 + \sum_{k=1}^{\infty} \binom{p}{k} a^{-k} M_k^k(f) \right)^{1/p} \\
&= a + a \sum_{j=1}^{\infty} \binom{1/p}{j} \left(\sum_{k=1}^{\infty} \binom{p}{k} a^{-k} M_k^k(f) \right)^j.
\end{aligned}$$

In the case $p = 0$, a similar calculation yields

$$M_0(f + ae_0) = aM(a^{-1}f + e_0) = a \exp \left(\sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{ja^j} M_j^j(f) \right).$$

2. Proofs

Proof of Theorem 1. First we consider the case $p \neq 0$. For $a > \max_{1 \leq i \leq n} |x_i|$, we have

$$\begin{aligned}
M_p(\mathbf{x} + \mathbf{a}) &= \left(\sum_{i=1}^n w_i (x_i + a)^p \right)^{1/p} = a \left(\sum_{i=1}^n w_i (1 + x_i/a)^p \right)^{1/p} \\
&= a \left(\sum_{i=1}^n w_i \sum_{k=0}^{\infty} \binom{p}{k} \left(\frac{x_i}{a} \right)^k \right)^{1/p} = a \left(\sum_{k=0}^{\infty} \binom{p}{k} \sum_{i=1}^n w_i \left(\frac{x_i}{a} \right)^k \right)^{1/p} \\
&= a \left(\sum_{k=0}^{\infty} \binom{p}{k} a^{-k} M_k^k(\mathbf{x}) \right)^{1/p} = a \left(1 + \sum_{k=1}^{\infty} \binom{p}{k} a^{-k} M_k^k(\mathbf{x}) \right)^{1/p} \\
&= a + a \sum_{j=1}^{\infty} \binom{1/p}{j} \left(\sum_{k=1}^{\infty} \binom{p}{k} a^{-k} M_k^k(\mathbf{x}) \right)^j
\end{aligned}$$

since $\sup_k M_k(\mathbf{x}) \leq \max_{1 \leq i \leq n} |x_i|$. From Eq. (3) we deduce

$$\left(\sum_{k=1}^{\infty} \binom{p}{k} \frac{M_k^k(\mathbf{x})}{a^k} \right)^j = j! \sum_{m=j}^{\infty} B_{m,j} \left[i! \binom{p}{i} M_i^i(\mathbf{x}) \right] \frac{a^{-m}}{m!}$$

($j = 1, 2, \dots$). Therefore, we conclude

$$M_p(\mathbf{x} + \mathbf{a}) = a + \sum_{m=1}^{\infty} \frac{1}{m! a^{m-1}} \sum_{j=1}^m j! \binom{1/p}{j} B_{m,j} \left[i! \binom{p}{i} M_i^i(\mathbf{x}) \right].$$

In the case $p = 0$ we have, for $a > \max_{1 \leq i \leq n} |x_i|$,

$$\begin{aligned} M_0(\mathbf{x} + \mathbf{a}) &= a \prod_{i=1}^n \left(1 + \frac{x_i}{a}\right)^{w_i} = a \exp\left(\sum_{i=1}^n w_i \log(1 + x_i/a)\right) \\ &= a \exp\left(\sum_{i=1}^n w_i \sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j} \left(\frac{x_i}{a}\right)^j\right) \\ &= a \exp\left(\sum_{j=1}^{\infty} \frac{(-1)^{j+1}}{j a^j} M_j^j(\mathbf{x})\right). \end{aligned}$$

With (4) we conclude that

$$M_0(\mathbf{x} + \mathbf{a}) = a + \sum_{m=1}^{\infty} \frac{(-1)^m}{m! a^{m-1}} Y_m[-(i-1)! M_i^i(\mathbf{x})].$$

This completes the proof of Theorem 1. \square

Proof of Theorem 5. With regard to Remark 2 application of Theorem 1 yields the asymptotic formula

$$\begin{aligned} M_p(\mathbf{a} + t\mathbf{x}) - M_q(\mathbf{a} + t\mathbf{x}) &= \frac{t^2}{a} \frac{p-q}{2} (M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})) \\ &\quad + \frac{t^2}{a} \sum_{k=1}^{\infty} \frac{(t/a)^k}{(k+1)!} \sum_{j=1}^{k+1} j! \left[\binom{1/p}{j} \mathbf{B}_{k+1,j} \left[i! \binom{p}{i} M_i^i(\mathbf{x}) \right] \right. \\ &\quad \left. - \binom{1/q}{j} \mathbf{B}_{k+1,j} \left[i! \binom{q}{i} M_i^i(\mathbf{x}) \right] \right]. \end{aligned}$$

Using the analogous relation for $M_r(\mathbf{a} + t\mathbf{x}) - M_s(\mathbf{a} + t\mathbf{x})$ the complete asymptotic expansion follows. The special case

$$\begin{aligned} &M_p(\mathbf{a} + t\mathbf{x}) - M_q(\mathbf{a} + t\mathbf{x}) \\ &= \frac{p-q}{2a} \frac{t^2}{M_2^2(\mathbf{x}) - M_1^2(\mathbf{x})} \\ &\quad \times \left[1 + \frac{(p+q-3)M_3^3(\mathbf{x}) - 3(p+q-2)M_1(\mathbf{x})M_2^2(\mathbf{x}) + (2p+2q-3)M_1^3(\mathbf{x})}{3a(M_2^2(\mathbf{x}) - M_1^2(\mathbf{x}))} t \right. \\ &\quad \left. + O(t^2) \right] \end{aligned}$$

as $t \rightarrow 0$ leads after some calculations to the explicit expressions of the first two coefficients. \square

References

- [1] M. Bjelica, Asymptotic linearity of mean values, *Math. Vesnik* 51 (1–2) (1999) 15–19.
- [2] L. Comtet, *Advanced Combinatorics*, Reidel, Dordrecht, 1974.
- [3] Bai-Ni Guo, Feng Qi, Inequalities for generalized weighted mean values of convex function, *Math. Inequal. Appl.* 4 (2) (2001) 195–202.
- [4] W. Rudin, *Real and Complex Analysis*, McGraw–Hill, 1974.